

PRIME IDEALS AND REGULAR SEQUENCES OF SYMMETRIC POLYNOMIALS

NEERAJ KUMAR

ABSTRACT. Let $S = \mathbb{C}[x_1, \dots, x_n]$ be a polynomial ring. Denote by p_a the power sum symmetric polynomial $x_1^a + \dots + x_n^a$. We consider the following two questions: Describe the subsets $A \subset \mathbb{N}$ such that the set of polynomials p_a with $a \in A$ generate a prime ideal in S or the set of polynomials p_a with $a \in A$ is a regular sequence in S . We produce a large families of prime ideals by exploiting Serre's criterion for normality [4, Theorem 18.15] with the help of arithmetic considerations, vanishing sums of roots of unity [9]. We also deduce several other results concerning regular sequences of symmetric polynomials.

1. INTRODUCTION

Let $S = \mathbb{C}[x_1, \dots, x_n]$ be a polynomial ring. A sequence of elements y_1, y_2, \dots, y_d in a ring S is called *regular sequence* on S if the ideal $\langle y_1, y_2, \dots, y_d \rangle$ is proper and for each i , the image of y_{i+1} is a nonzero divisor in $S/\langle y_1, \dots, y_i \rangle$.

Following the notation of Macdonald [10], let p_a, h_a and e_a denote the power sum symmetric polynomial, complete symmetric polynomial, and the elementary symmetric polynomial of degree a in S respectively. Let \mathbb{N} be the set of positive integers. For a given set $A \subset \mathbb{N}$, we denote by the set of power sum symmetric polynomials as $p_A = \{p_a \mid a \in A\}$. In this paper, we discuss the following two questions:

Question 1.1. *Let $S = \mathbb{C}[x_1, \dots, x_n]$ be a polynomial ring. For which subsets $A \subset \mathbb{N}$, the ideal generated by the set of polynomials p_A is a prime ideal in S .*

Question 1.2. *Let $S = \mathbb{C}[x_1, \dots, x_n]$ be a polynomial ring. For which subsets $A \subset \mathbb{N}$, the set of polynomials p_A is a regular sequence in S .*

Similarly, we ask these questions for the complete symmetric polynomials and the elementary symmetric polynomials. For obvious reasons, whenever p_A is a regular sequence generating a prime ideal and $p_b \notin \langle p_A \rangle$, then p_A, p_b is a regular sequence in S as well. More specifically, we will focus on Question 1.1.

These problems are fundamental in nature, and interesting, both in algebraic and geometric point of view. In a more general setting of Question 1.1, it is mentioned by Eisenbud that “In general it is extremely difficult to prove that a given ideal of polynomial is prime” [4, Chapter 10: pg.241]. The most powerful methods known for showing primeness of ideal are Hochster's method of “*principal radical system*” [7] and *Serre's criterion for normality* [4, Theorem 18.15]. We will use the Serre's criterion for normality in this paper.

The study of Question 1.2 began in the paper [1] in the dimension zero case by Conca, Krattenthaler, and Watanabe. The Question 1.2 is highly non-trivial for

Key words: Regular sequence, Prime ideal, Symmetric polynomial.

2010 Mathematics Subject Classification: Primary 13A15, Secondary 14M10, 12E05.

$n \geq 3$. For $n = 3$, a beautiful conjecture of Conca, Krattenthaler, and Watanabe states that given a positive integers $a < b < c$ with $\gcd(a, b, c) = 1$, p_a, p_b, p_c form a regular sequence in $\mathbb{C}[x_1, x_2, x_3]$ if and only if $abc \equiv 0 \pmod{6}$, see [1, Conjecture 2.15]. The necessary condition follows from [1, Lemma 2.8]. For partial evidence in support of sufficient condition, see [1, Theorem 2.11]. Similarly the authors, also formulated a conjecture, when three complete symmetric polynomials form a regular sequence in $\mathbb{C}[x_1, x_2, x_3]$, see [1, Conjecture 2.17]. In a joint paper with Martino [8], we could provide evidence for these conjectures by proving it in special instances. Then we employed the technique of Serre's criterion to show the primeness of an ideal. For instance, we have shown that the ideal $I = \langle p_a, p_{a+1}, \dots, p_{a+m-1} \rangle$ is prime in S if $m < n - 1$, see [8, Theorem 3.3]. We have also shown that the ideal $I = \langle p_1, p_{2m} \rangle$, where $m \in \mathbb{N}$, is prime in $\mathbb{C}[x_1, \dots, x_4]$, see [8, Proposition 4.1]. In this way, we succeeded to give more families of regular sequences. With the help of Computer calculations, we proposed, Conjecture 4.5 and Conjecture 4.6 in [8]. One of the main result of this paper, Theorem 3.8, partially answers the Conjecture 4.6 in [8].

In this paper, we have managed to produce families of prime ideals by exploiting Serre's criterion with the help of arithmetic considerations, vanishing sums of roots of unity [9]. The main results of the paper are the following:

- (i) Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring with $n \geq 4$. Let $I = \langle p_a, p_b \rangle$, where $a, b \in \mathbb{N}$. Let $b - a = n_0$. Suppose q_1 is the smallest prime factor in the factorization of n_0 . If $q_1 > \max\{n, a\}$, then I is a prime ideal in S .
- (ii) Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring with $n \geq 3$. Let $a \in \mathbb{N}$ and $m < n - 1$. Let $I = \langle p_a, p_{2a}, \dots, p_{ma} \rangle$. Then I is a prime ideal in S .

Section 2 contains preliminary results. In Section 3, we discuss the problem of whether two power sum symmetric polynomials generate a prime ideal in S for $n \geq 4$. We answer this to certain extent purely in terms of arithmetic conditions of the degree of polynomials and the number of indeterminates, see Theorem 3.8. Let $I = \langle p_a, p_{2a}, \dots, p_{ma} \rangle$, where $a \in \mathbb{N}$, and $m < n - 1$. We show that I is a prime ideal in S for all $n \geq 3$, see Theorem 3.13. For $n \geq 3$, we show that $\frac{\partial h_a}{\partial x_1}, \dots, \frac{\partial h_a}{\partial x_n}$ form a regular sequence in S for all $a \geq 2$. We also show that any two complete symmetric polynomial form a regular sequence in S for all $n \geq 3$. Similar results also hold for the power sum and elementary symmetric polynomials, see Lemma 3.1 and Proposition 3.7. Computer calculations in CoCoA [2] suggest that $I = \langle h_1, h_{2m} \rangle$, where $m \in \mathbb{N}$, should be a prime ideal in $\mathbb{C}[x_1, \dots, x_4]$. It is obvious for $m = 1$. We provide evidence for $m = 2$ in the Example 3.16.

2. GENERALITIES AND PRELIMINARY RESULTS

Let $S = \mathbb{C}[x_1, \dots, x_n]$ be a polynomial ring. Let p_a , h_a and e_a be the power sum symmetric polynomial, complete symmetric polynomial, and the elementary symmetric polynomial of degree a in S respectively, that is,

$$\begin{aligned} p_a(x_1, x_2, \dots, x_n) &:= \sum_{i=1}^n x_i^a, \\ h_a(x_1, x_2, \dots, x_n) &:= \sum_{1 \leq i_1 < i_2 < \dots < i_a \leq n} x_{i_1} x_{i_2} \cdots x_{i_a}, \\ e_a(x_1, x_2, \dots, x_n) &:= \sum_{1 \leq i_1 < i_2 < \dots < i_a \leq n} x_{i_1} x_{i_2} \cdots x_{i_a}. \end{aligned}$$

For instance, for $n = 3$ and $a = 2$, one has

$$\begin{aligned} p_2(x_1, x_2, x_3) &= x_1^2 + x_2^2 + x_3^2, \\ h_2(x_1, x_2, x_3) &= x_1^2 + x_2^2 + x_3^2 + x_1x_2 + x_1x_3 + x_2x_3, \\ e_2(x_1, x_2, x_3) &= x_1x_2 + x_1x_3 + x_2x_3. \end{aligned}$$

For these symmetric polynomials, we have the following *Newton's formula*, see [10, Equation's 2.6', 2.11, 2.11' respectively]:

$$\sum_{i=0}^n (-1)^i e_i h_{n-i} = 0 \text{ for all } n \geq 1, \quad (1)$$

$$ah_a = \sum_{i=1}^a p_i h_{a-i} \text{ for all } a \geq 1, \quad (2)$$

$$ne_n = \sum_{i=1}^n (-1)^{i-1} e_{n-i} p_i \text{ for all } n \geq 1. \quad (3)$$

We will use the following lemma to prove the smoothness of symmetric polynomials h_a , e_a , and p_a in the Lemma 3.2. We will also use Lemma 2.1 in the Example 3.16.

Lemma 2.1. (Technical Lemma)

Let $S = \mathbb{C}[x_1, \dots, x_n]$ be a polynomial ring. Then for the symmetric polynomials h_a , e_a , and p_a , one has the following:

- (i) $\frac{\partial h_a}{\partial x_i} = h_{a-1} + x_i \frac{\partial h_{a-1}}{\partial x_i}$ and $\sum_{i=1}^n \frac{\partial h_a}{\partial x_i} = (n + a - 1)h_{a-1}$.
- (ii) $\frac{\partial e_a}{\partial x_i} = e_{a-1} - x_i \frac{\partial e_{a-1}}{\partial x_i}$ and $\sum_{i=1}^n \frac{\partial e_a}{\partial x_i} = (n - a + 1)e_{a-1}$.
- (iii) $\frac{\partial p_a}{\partial x_i} = ax_i^{a-1}$ and $\sum_{i=1}^n \frac{\partial p_a}{\partial x_i} = ap_{a-1}$.

Proof. We can write $h_a = x_i h_{a-1} + g$ for some polynomial g not involving x_i . Taking partial derivative of h_a w.r.t. x_i , one obtains

$$\frac{\partial h_a}{\partial x_i} = h_{a-1} + x_i \frac{\partial h_{a-1}}{\partial x_i}. \quad (4)$$

By Euler's formula, we have $\sum_{i=1}^n x_i \frac{\partial h_{a-1}}{\partial x_i} = (a-1)h_{a-1}$. Thus, we conclude that

$$\sum_{i=1}^n \frac{\partial h_a}{\partial x_i} = (n + a - 1)h_{a-1}.$$

Taking partial derivative of e_a w.r.t. x_i , one obtains

$$\frac{\partial e_a}{\partial x_i} = e_{a-1} - x_i \frac{\partial e_{a-1}}{\partial x_i}. \quad (5)$$

Proceeding as before, we conclude that

$$\sum_{i=1}^n \frac{\partial e_a}{\partial x_i} = (n - a + 1)e_{a-1}.$$

The claim of (iii) is obvious. \square

For a given natural number m , consider the m -th roots of unity in the field of complex number \mathbb{C} . We may ask ourselves for which natural numbers n and k , do there exist m -th roots of unity $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ such that $\alpha_1^k + \alpha_2^k + \dots + \alpha_n^k = 0$? We define such an equation to be *Vanishing sum of k -th power of m -th roots of unity of weight n* . For $k = 1$, such an equation is said to be a *vanishing sum of m -th roots of unity of weight n* . For instance, for $m = 10$ and $k = 1$, the set of n 's is $\{0, 2, 4, 5, 6, 7, 8, 9, 10, \dots\}$. For a given m and k , let $W(m, k)$ be the set of weights n for which there exists a vanishing sum $\alpha_1^k + \alpha_2^k + \dots + \alpha_n^k = 0$, where α_i is a m -th roots of unity. If m has a prime factorization of the form $q_1^{a_1} q_2^{a_2} \dots q_r^{a_r}$, then by the theorem of Lam and Leung [9, Theorem 5.2], the weight set $W(m, 1)$ is exactly given by $\mathbb{N}q_1 + \dots + \mathbb{N}q_r$. Poonen and Rubinstein [13] have classified all minimal vanishing sums $\alpha_1 + \dots + \alpha_n = 0$ of weight $n \leq 12$. For similar treatment by Mann, one may also see [11].

Remark 2.2. Note that by [9, Theorem 5.2], for a given $m \in \mathbb{N}$, $W(m, 1)$ depends only on the prime divisors of m , and not on the multiplicity to which they occur in the factorization of m . The theorem also shows that any (non-empty) vanishing sum of m -th roots of unity must have weight $\geq q_1$, where q_1 is the smallest prime divisor of m .

Lemma 2.3. *Let $m \in \mathbb{N}$ and $\alpha_1, \dots, \alpha_n$ be a subset of m -th roots of unity in \mathbb{C} . Suppose that q_1 is the smallest prime factor in the factorization of m . Let $k \in \mathbb{N}$ be any natural number such that $q_1 > \max\{n, k\}$. Then one has $\alpha_1^k + \alpha_2^k + \dots + \alpha_n^k \neq 0$.*

Proof. The claim follows from Remark 2.2. \square

3. PRIME IDEALS AND REGULAR SEQUENCES

We begin this section with a useful lemma, which states that all the partial derivatives of a complete symmetric polynomial form a regular sequence in the polynomial ring, see Lemma 3.1. Then we use Lemma 3.1 to show that the complete symmetric polynomials and their partial derivatives are smooth polynomials, see Lemma 3.2 and Lemma 3.3. Then we recall the irreducibility of Schur polynomial in the polynomial ring, see [3, Theorem 3.1]. In a special case, we discuss the smoothness of Schur polynomial, see Example 3.5. Then we discuss the main results of this paper, Theorem 3.8 and Theorem 3.13 respectively. We also discuss the case of two complete symmetric polynomials generating a prime ideal in the polynomial ring.

Lemma 3.1. *Let $n \in \mathbb{N}$ with $n \geq 3$. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring. Let $a \geq 2$. Then the following holds:*

- (i) $\frac{\partial h_a}{\partial x_1}, \dots, \frac{\partial h_a}{\partial x_n}$ form a regular sequence in S .
- (ii) $\frac{\partial p_a}{\partial x_1}, \dots, \frac{\partial p_a}{\partial x_n}$ form a regular sequence in S .
- (iii) $\frac{\partial e_a}{\partial x_1}, \dots, \frac{\partial e_a}{\partial x_n}$ form a regular sequence in S for all $a < n$.

Proof. Let the ideal generated by all the partial derivatives of h_a be J_a , that is,

$$J_a = \left\langle \frac{\partial h_a}{\partial x_1}, \dots, \frac{\partial h_a}{\partial x_n} \right\rangle. \quad (6)$$

We have $ah_a = \sum x_i \frac{\partial h_a}{\partial x_i}$. Thus, clearly $h_a \in J_a$. By Lemma 2.1 (i), we have

$$(n+a-1)h_{a-1} = \sum_{i=1}^n \frac{\partial h_a}{\partial x_i} \in J_a.$$

That is, $h_{a-1} \in J_a$. By Lemma 2.1 (i), we have

$$\frac{\partial h_{a+1}}{\partial x_i} = h_a + x_i \frac{\partial h_a}{\partial x_i} \in J_a.$$

Thus, we observe that $J_{a+1} \subset J_a$. Moreover, $h_{a+1} \in J_{a+1}$. Proceeding similarly, we have a chain of containment of ideals

$$J_{a+n-2} \subset \cdots \subset J_{a+1} \subset J_a.$$

Therefore, we have $h_{a-1}, h_a, h_{a+1}, \dots \subset h_{a+n-2} \in J_a$. Recall that by [1, Proposition 2.9], any n consecutive complete symmetric polynomials $h_{a-1}, h_a, \dots, h_{a+n-2}$ form a regular sequence in S for all $a \geq 2$. Thus, we conclude that $\text{ht}(J_a) = n$, moreover J_a is a complete intersection ideal in S . The claim (ii) is obvious. Let the ideal generated by all the partial derivatives of e_a be E_a . The proof of (iii) is similar to above, except the fact that this time, we choose carefully a 's such that $e_a \in E_a$. The reason for this is that $e_a = 0$ for all $a > n$. We want to use the fact that $e_1, e_2, \dots, e_n \in E_a$. As we know that e_1, \dots, e_n form a regular sequence in S . \square

In the following lemma, we discuss the smoothness of the symmetric polynomials h_a , e_a and p_a .

Lemma 3.2. *Let $n \in \mathbb{N}$ with $n \geq 3$. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring. Then the following holds:*

- (i) h_a is smooth, hence an irreducible element in S for all $a \geq 1$.
- (ii) e_a is smooth, hence an irreducible element in S for all $1 \leq a \leq n-1$.
- (iii) p_a is smooth, hence an irreducible element in S for all $a \geq 1$.

Proof. If $a = 1$, the claims are obvious. We will give an elementary proof of (i). The proof of (ii) and (iii) are similar.

Proof of (i): If $h_a = f \cdot g$ with f and g non constant polynomial, then f and g have to be homogeneous. By Bezout theorem, the hypersurfaces $f = 0$ and $g = 0$ intersects in the projective space \mathbb{P}^{n-1} , since $n \geq 3$. This gives a singular point on the hypersurface $h_a = 0$. So, it suffices to prove that $h_a, \frac{\partial h_a}{\partial x_1}, \dots, \frac{\partial h_a}{\partial x_n}$ have no common zero in $\mathbb{C}^n - \{0\}$. This claim follows from Lemma 3.1 (i). \square

In the following lemma, we discuss the smoothness of the partial derivatives of the complete symmetric polynomials.

Lemma 3.3. *Let $n \in \mathbb{N}$ with $n \geq 3$. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring. Let $a \geq 3$. Then $\frac{\partial h_a}{\partial x_i}$ is smooth, and hence irreducible in S .*

Proof. It is enough to show that $\frac{\partial h_a}{\partial x_1}$ is smooth. Similar to the proof given in Lemma 3.2, it suffices to show that all its partial derivatives $\frac{\partial^2 h_a}{\partial x_1 \partial x_1}, \frac{\partial^2 h_a}{\partial x_1 \partial x_2}, \dots, \frac{\partial^2 h_a}{\partial x_1 \partial x_n}$ have no common zero in $\mathbb{C}^n - \{0\}$. We set few notations for convenience, for instance, let $g_{a,i} = \frac{\partial h_a}{\partial x_i}$. Let the ideal generated by all the partial derivatives of $g_{a,1}$ be I_a :

$$I_a = \left\langle \frac{\partial g_{a,1}}{\partial x_1}, \frac{\partial g_{a,1}}{\partial x_2}, \dots, \frac{\partial g_{a,1}}{\partial x_n} \right\rangle.$$

Also note that $\frac{\partial g_{a,j}}{\partial x_i} = \frac{\partial g_{a,i}}{\partial x_j}$. With this observation, one obtains

$$I_a = \left\langle \frac{\partial g_{a,1}}{\partial x_1}, \frac{\partial g_{a,2}}{\partial x_1}, \dots, \frac{\partial g_{a,n}}{\partial x_1} \right\rangle.$$

Taking partial derivatives of $g_{a,i}$ w.r.t. x_1 , for $i = 1, \dots, n$,

$$\begin{aligned} \frac{\partial g_{a,1}}{\partial x_1} &= \frac{\partial h_{a-1}}{\partial x_1} + x_1 \frac{\partial^2 h_a}{\partial x_1 \partial x_1} + \frac{\partial h_{a-1}}{\partial x_1}, \\ \frac{\partial g_{a,2}}{\partial x_1} &= \frac{\partial h_{a-1}}{\partial x_1} + x_2 \frac{\partial^2 h_a}{\partial x_2 \partial x_1} \\ &\vdots \\ \frac{\partial g_{a,n}}{\partial x_1} &= \frac{\partial h_{a-1}}{\partial x_1} + x_n \frac{\partial^2 h_a}{\partial x_n \partial x_1} \end{aligned}$$

Then summing up, we get

$$\begin{aligned} \sum_{i=1}^n \frac{\partial g_{a,i}}{\partial x_1} &= (n+1) \frac{\partial h_{a-1}}{\partial x_1} + \sum_{i=1}^n x_i \frac{\partial}{\partial x_i} \left(\frac{\partial h_{a-1}}{\partial x_1} \right) \\ &= (\sharp) \frac{\partial h_{a-1}}{\partial x_1}, \end{aligned}$$

for some integer number \sharp , which is irrelevant. Thus, we conclude that $\frac{\partial h_{a-1}}{\partial x_1} \in I_a$. Also $g_{a,1} \in I_a$. We also know that $g_{a,1} = \frac{\partial h_a}{\partial x_1} = h_{a-1} + x_1 \frac{\partial h_{a-1}}{\partial x_1}$. Thus, we get $h_{a-1} \in I_a$. Now consider the ideal

$$I_{a+1} = \left\langle \frac{\partial g_{a+1,1}}{\partial x_1}, \frac{\partial g_{a+1,1}}{\partial x_2}, \dots, \frac{\partial g_{a+1,1}}{\partial x_n} \right\rangle.$$

Proceeding similarly, we obtain

$$\begin{aligned} \sum_{i=1}^n \frac{\partial g_{a+1,i}}{\partial x_1} &= (n+1) \frac{\partial h_a}{\partial x_1} + \sum_{i=1}^n x_i \frac{\partial}{\partial x_i} \left(\frac{\partial h_a}{\partial x_1} \right) \\ &= (\sharp_1) \frac{\partial h_a}{\partial x_1}, \end{aligned}$$

for some integer number \sharp_1 , which is irrelevant. Thus, we conclude that $\frac{\partial h_a}{\partial x_1} \in I_{a+1}$. Also $g_{a+1,1} \in I_{a+1}$. We also know that $g_{a+1,1} = \frac{\partial h_{a+1}}{\partial x_1} = h_a + x_1 \frac{\partial h_a}{\partial x_1}$. Thus, we get $h_a \in I_{a+1}$. Also note that $\frac{\partial g_{a+1,1}}{\partial x_i} \in I_a$ for all i . That is, $I_{a+1} \subset I_a$. Proceeding in a similar way, we obtain the following relations

$$I_{a+j+1} \subset I_{a+j}, \text{ and } h_{a+j-1} \in I_{a+j}$$

for all $j \geq 0$. Thus, using similar argument as in the Lemma 3.1, we conclude that $\text{ht}(I_a) = n$, moreover I_a is a complete intersection ideal in S . Thus the claim follows. \square

Following Macdonald [10], the *Schur polynomial* is defined as

$$s_\lambda = s_\lambda(x_1, \dots, x_n) = \frac{\det(x_i^{\lambda_j + n - j})_{1 \leq i, j \leq n}}{\det(x_i^{n-j})_{1 \leq i < j \leq n}},$$

where $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ is the partition of non-negative integers with $\lambda_i \geq \lambda_{i+1}$ for $i = 1, \dots, n-1$. For Schur polynomial, we have the following relation, see [10, Equation 3.4]:

$$s_\lambda = \det(h_{\lambda_i - i + j})_{1 \leq i, j \leq n} \quad (7)$$

where $n \geq l(\lambda)$.

Remark 3.4. Irreducibility of Schur polynomial is discussed by Dvornicich and Zannier in [3]. Let $n \geq 3$. For a given partition $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$, where $\lambda_1 > \lambda_2 > \dots > \lambda_n$ with $\lambda_n = 0$ and $\gcd(\lambda_1, \lambda_2, \dots, \lambda_{n-1}) = 1$, then the Schur polynomial $s_\lambda(x_1, \dots, x_n)$ is irreducible in $S = \mathbb{C}[x_1, x_2, \dots, x_n]$, see [3, Theorem 3.1]. Recall that h_a and e_a are special forms of a Schur polynomial. Simply note that the irreducibility of h_a and e_a does not follow from [3, Theorem 3.1]. In the Lemma 3.2, we not only show the irreducibility of these polynomials, but also the smoothness. The obtained results partially extend the domain of partition for the irreducibility of Schur polynomial.

Assuming that the Schur polynomial $s_\lambda(x_1, \dots, x_n)$ is irreducible in S . One may ask, whether is it true that the Schur polynomial is also smooth? The answer is positive in the case of h_a and e_a . Computational evidence shows that the answer is negative in general. However, in a special case, when the partition λ is of the form $(\lambda_1, 1, 0)$ for $\lambda_1 \geq 2$, then the Schur polynomial $s_\lambda(x_1, x_2, x_3)$ turns out to be smooth in $\mathbb{C}[x_1, x_2, x_3]$. We discuss the proof of this in the following example:

Example 3.5. Let $S = \mathbb{C}[x_1, x_2, x_3]$ be a polynomial ring. Let $\lambda = (\lambda_1, 1, 0)$ be the partition. Then the Schur polynomial $s_\lambda(x_1, x_2, x_3)$ is smooth in S for all $\lambda_1 \geq 2$.

Proof. By (7), we have

$$s_\lambda = s_{(\lambda_1, 1, 0)} = h_1 h_{\lambda_1} - h_{\lambda_1 + 1}.$$

Similar to the proof given in Lemma 3.2, it suffices to show that all its partial derivatives $\frac{\partial s_\lambda}{\partial x_1}, \frac{\partial s_\lambda}{\partial x_2}, \frac{\partial s_\lambda}{\partial x_3}$ have no common zero in $\mathbb{C}^3 - \{0\}$. Taking partial derivatives of s_λ w.r.t. x_i , we get

$$\begin{aligned} \frac{\partial s_\lambda}{\partial x_i} &= h_{\lambda_1} + h_1 \frac{\partial h_{\lambda_1}}{\partial x_i} - \frac{\partial h_{\lambda_1 + 1}}{\partial x_i} \\ &= (h_1 - x_i) \frac{\partial h_{\lambda_1}}{\partial x_i} \text{ for all } i = 1, 2, 3. \end{aligned}$$

We see that $h_1 \neq x_i$, unless $x_i = 0$ for all i . Thus the common zero of $\frac{\partial h_{\lambda_1}}{\partial x_1}, \frac{\partial h_{\lambda_1}}{\partial x_2}, \frac{\partial h_{\lambda_1}}{\partial x_3}$ is also a common zero of $\frac{\partial s_\lambda}{\partial x_1}, \frac{\partial s_\lambda}{\partial x_2}, \frac{\partial s_\lambda}{\partial x_3}$. By Lemma 3.1, we conclude that $\langle \frac{\partial s_\lambda}{\partial x_1}, \frac{\partial s_\lambda}{\partial x_2}, \frac{\partial s_\lambda}{\partial x_3} \rangle$ is a complete intersection ideal in S . Thus the claim follows. \square

We record the following convention, which we will follow from here onwards throughout this paper:

Conventions 3.6. When we list the symmetric polynomials $f_{i_1}, f_{i_2}, \dots, f_{i_k}$ with respective degrees $\deg(f_{i_j}) = i_j$, we always assume that $i_1 < i_2 < \dots < i_k$, unless otherwise specified.

In the following proposition, we will see that any two complete symmetric polynomials always form a regular sequence in $S = \mathbb{C}[x_1, \dots, x_n]$ for $n \geq 3$. Similar results also hold for the power sum and elementary symmetric polynomials.

Proposition 3.7. *Let $n \in \mathbb{N}$ with $n \geq 3$. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring. Then the following holds:*

- (i) h_a, h_b form a regular sequence.
- (ii) e_a, e_b form a regular sequence for all $1 \leq a < b \leq n-1$.
- (iii) p_a, p_b form a regular sequence.

Proof. We will prove (i). By Lemma 3.2, h_a is an irreducible polynomial in S . Hence $S/\langle h_a \rangle$ is a domain. Now h_b being an irreducible polynomial in S , can not be factored into lower degree complete symmetric polynomials h_a . So, h_b is a nonzero divisor in $S/\langle h_a \rangle$ for $b > a$. Hence h_a, h_b form a regular sequence in S . Proof of (ii) and (iii) are similar. \square

We have seen that any two power sum polynomials p_a, p_b form a regular sequence in $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ for $n \geq 3$. We would like to know when two power sum polynomials generate a prime ideal in S for $n \geq 4$. In a special case, some answers are known due to [8, Theorem 4.3 and Proposition 4.3]. In the following theorem, we will answer this to certain extent, when two power sum polynomials generate a prime ideal in S for $n \geq 4$, purely in terms of arithmetic conditions of the degree of polynomials and the number of indeterminates.

Theorem 3.8. *Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring with $n \geq 4$. Let $I = \langle p_a, p_b \rangle$, where $a, b \in \mathbb{N}$. Let $b - a = n_0$. Suppose q_1 is the smallest prime factor in the factorization of n_0 . If $q_1 > \max\{n, a\}$, then I is a prime ideal in S . Moreover p_a, p_b, p_c form a regular sequence in S for all $p_c \notin \langle p_a, p_b \rangle$.*

Proof. If $n_0 = 1$, then I is a prime ideal in S follows from [8, Theorem 4.3]. Thus, we assume that $n_0 \geq 2$. Let $R = S/I$. We compute the Jacobian of I up to scalar (We can ignore the coefficients, since we are in the field of characteristic zero.), say Jacobian is J :

$$J = \begin{pmatrix} x_1^{a-1} & x_2^{a-1} & \dots & x_n^{a-1} \\ x_1^{b-1} & x_2^{b-1} & \dots & x_n^{b-1} \end{pmatrix}.$$

Let $J' = I_2(J)$, denotes the ideal generated by 2×2 minors of Jacobian. Also $\text{ht}(I) = 2$, since I is generated by a regular sequence of length 2. The determinants of 2×2 minors of the Jacobian can be written as

$$J' = \langle x_j^{a-1} x_i^{a-1} (x_j^{b-a} - x_i^{b-a}) \rangle \text{ for } 1 \leq i < j \leq n.$$

Thus, we have

$$I + J' = \langle p_a, p_b, x_j^{a-1} x_i^{a-1} (x_j^{b-a} - x_i^{b-a}) \rangle \text{ for } 1 \leq i < j \leq n.$$

Claim: $\sqrt{I + J'} = (x_1, x_2, \dots, x_n)$.

Suppose not, that is, there exists $w = (w_1, w_2, \dots, w_n) \in \mathbb{P}^{n-1}$ with $w \in Z(I + J')$. Since w is in \mathbb{P}^{n-1} , we may assume $w = (1, y_1, y_2, \dots, y_{n-1})$. As $w \in Z(I + J')$, we have that $y_i^{b-a} = y_i^{n_0} = 1$ for all $i = 1, \dots, n-1$, moreover w also satisfies p_a, p_b . Therefore we have

$$1 + y_1^a + y_2^a + \dots + y_{n-1}^a = 0 \text{ and } 1 + y_1^b + y_2^b + \dots + y_{n-1}^b = 0.$$

Both the equations reduce to the existence of solution of $1 + y_1^a + y_2^a + \dots + y_{n-1}^a = 0$. We use the fact that all the y_i 's are n_0 -th roots of unity, say $1, \zeta_1, \dots, \zeta_{n_0-1}$. Suppose q_1 is the smallest prime factor in the factorization of n_0 . If $q_1 > \max\{n, a\}$, then

it follows from Lemma 2.3 that $1 + y_1^a + y_2^a + \cdots + y_{n-1}^a \neq 0$. So, the only possible solution has to be the trivial solution. Hence the claim is proved.

Thus $\text{ht}(I + J') = n$ and $\dim S/(I + J') = 0$. The co-dimension of J' in S is $n - 2$. By [4, Theorem 18.15], R is a product of normal domain, since $n \geq 4$. Thus, we can write $R = R_1 \times \cdots \times R_k$. Since R is a standard graded \mathbb{C} -algebra with $R_0 = \mathbb{C}$, also $R_0 = (R_1)_0 \times \cdots \times (R_k)_0 = \mathbb{C}^k$. Hence $k = 1$. Thus R is a normal domain and I is a prime ideal in S . \square

Remark 3.9. Let $p_1, p_2, p_5 \in \mathbb{C}[x_1, \dots, x_4]$. By Newton's formula (3), we observe that $p_5 \equiv 0 \pmod{(p_1, p_2)}$. Hence $p_5 \in \langle p_1, p_2 \rangle$. Replacing x_i by x_i^d , we may also conclude that $p_{5d} \in \langle p_d, p_{2d} \rangle$. Thus in the hypothesis of Theorem 3.8, we need the condition $p_c \notin \langle p_a, p_b \rangle$.

Remark 3.10. Computer calculations in CoCoA [2] suggest that whenever $I = \langle p_a, p_b \rangle$ is a prime ideal in $\mathbb{C}[x_1, x_2, x_3, x_4]$, then p_a, p_b, p_c form a regular sequence, except p_a, p_{2a}, p_{5a} . It is clear from Remark 3.9 that p_a, p_{2a}, p_{5a} do not form a regular sequence. If this computational claim can be answered, then it will prove [8, Conjecture 4.5] to certain extent.

Proposition 3.11. *Let I and J be the prime ideal in $K[x_1, \dots, x_n]$ and $K[y_1, \dots, y_m]$ respectively, where K is an algebraically closed field. Let (I, J) be the ideal generated by elements of I and J in $K[x_1, \dots, x_n, y_1, \dots, y_m]$. Then (I, J) is a prime ideal in $K[x_1, \dots, x_n, y_1, \dots, y_m]$.*

Proof. There is a standard isomorphism

$$K[x_1, \dots, x_n]/I \otimes K[y_1, \dots, y_m]/J \cong K[x_1, \dots, x_n, y_1, \dots, y_m]/(I, J)$$

by sending $f \otimes g \mapsto fg$. We see that both $K[x_1, \dots, x_n]/I$ and $K[y_1, \dots, y_m]/J$ are integral domains as well as K -algebras. By [12, Proposition 4.15(b)], the tensor product of $K[x_1, \dots, x_n]/I$ and $K[y_1, \dots, y_m]/J$ is also an integral domain, since K is algebraically closed field. Hence the claim follows. \square

Remark 3.12. Note that the goal of Proposition 3.11 is to generate more families of prime ideals from given prime ideals.

By [1, Proposition 2.9], we know that p_1, p_2, \dots, p_n form a regular sequence in $S = \mathbb{C}[x_1, x_2, \dots, x_n]$. We also know that a subset of a regular sequence is again a regular sequence. Thus p_1, p_2, \dots, p_m also form a regular sequence for all $m < n$. Then by [1, Lemma 2.2], we conclude that $p_a, p_{2a}, \dots, p_{ma}$ also form a regular sequence. Let $I = \langle p_a, p_{2a}, \dots, p_{ma} \rangle$, where $a \in \mathbb{N}$, and $m < n - 1$. Let $R = S/I$. Then R is a Cohen-Macaulay ring. In the following theorem, we will show that I is a prime ideal in S . We show this by proving that R is a normal domain using Serre's criterion for normality.

Theorem 3.13. *Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring with $n \geq 3$. Let $a \in \mathbb{N}$. Let $I = \langle p_a, p_{2a}, \dots, p_{ma} \rangle$, where $m < n - 1$. Then I is a prime ideal in S .*

Proof. For $a = 1$, it follows from [8, Proposition 4.3]. Assume $a > 1$. Let $R = S/I$. We compute the Jacobian of I up to scalar, say Jacobian is J :

$$J = \begin{pmatrix} x_1^{a-1} & x_2^{a-1} & \cdots & x_n^{a-1} \\ x_1^{2a-1} & x_2^{2a-1} & \cdots & x_n^{2a-1} \\ \vdots & \vdots & \cdots & \vdots \\ x_1^{ma-1} & x_2^{ma-1} & \cdots & x_n^{ma-1} \end{pmatrix}.$$

We can ignore the coefficients, since we are in the field of characteristic zero. We have $\text{ht}(I) = m$, since I is generated by a regular sequence of length m . Let $J' = I_m(J)$, denotes the ideal generated by $m \times m$ minors of Jacobian J . The determinants of $m \times m$ minors of the Jacobian can be written as

$$J' = \langle x_{i_1}^{j_1} x_{i_2}^{j_2} \cdots x_{i_m}^{j_m} \prod_{1 \leq a < b \leq m} (x_{i_a}^{j_a} - x_{i_b}^{j_b}) \rangle \text{ for } 1 \leq i_1 < i_2 < \cdots < i_m \leq n,$$

where j_1, j_2, \dots, j_m are some positive integers. Therefore

$$I + J' = \langle p_a, p_{2a}, \dots, p_{ma}, x_{i_1}^{j_1} x_{i_2}^{j_2} \cdots x_{i_m}^{j_m} \prod_{1 \leq a < b \leq m} (x_{i_a}^{j_a} - x_{i_b}^{j_b}) \rangle.$$

Claim: $\sqrt{I + J'} = (x_1, x_2, \dots, x_n)$.

Suppose not, that is, there exists $w \in \mathbb{P}^{n-1}$ with $w \in Z(I + J')$. Then the vector w can have at the most $m - 1$ distinct nonzero coordinates. If w has m or more than m distinct nonzero coordinates, then $w \notin Z(J')$. Say w has v distinct nonzero coordinates. We can write

$$w = (w_1, \dots, w_1, w_2, \dots, w_2, \dots, w_v, \dots, w_v, 0, 0, \dots, 0),$$

where w_i appears β_i times and $v \leq m - 1$. Also w should satisfy p_{ia} for $i = 1, \dots, m$. Thus, we have

$$\beta_1 w_1^{ia} + \beta_2 w_2^{ia} + \cdots + \beta_v w_v^{ia} = 0 \text{ for } i = 1, 2, \dots, m.$$

This is a system of equation, which can be represented in the matrix form with m rows and v columns as

$$\begin{pmatrix} 1 & 1 & \cdots & 1 \\ w_1^a & w_2^a & \cdots & w_v^a \\ \vdots & \vdots & \cdots & \vdots \\ w_1^{(m-1)a} & w_2^{(m-1)a} & \cdots & w_v^{(m-1)a} \end{pmatrix} \begin{pmatrix} \beta_1 w_1^a \\ \beta_2 w_2^a \\ \vdots \\ \beta_v w_v^a \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}.$$

We know that neither $\beta_i = 0$ nor $w_i = 0$ for $i = 1, \dots, v$. So $\beta_i w_i^a \neq 0$ for $i = 1, \dots, v$. We can choose the matrix say M with first v rows out of m rows and look for the solution. The matrix M is of full rank since $w_i \neq w_j$ for $i \neq j$, so the only possible solution has to be the trivial solution. Hence the claim is proved. By similar argument as used in Theorem 3.8, we conclude that R is a normal domain and I is a prime ideal in S . \square

For $n \geq 4$, we know that the ideal $\langle p_1, p_2, \dots, p_m \rangle$ is prime in $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ for all $m < n - 1$, see [8, Theorem 4.3]. We will see in the following theorem that similar result holds for the complete symmetric polynomials and the elementary symmetric polynomials.

Proposition 3.14. *Let p_a, h_a , and e_a in the polynomial ring $S = \mathbb{C}[x_1, x_2, \dots, x_n]$, with $n \geq 4$. Let $m < n - 1$. Then one has*

$$\langle p_1, p_2, \dots, p_m \rangle = \langle h_1, h_2, \dots, h_m \rangle = \langle e_1, e_2, \dots, e_m \rangle.$$

Therefore the ideals $\langle h_1, h_2, \dots, h_m \rangle$ and $\langle e_1, e_2, \dots, e_m \rangle$ are also prime in S .

Proof. It follows from simple observation in the ring of symmetric polynomials that the algebra generated by the power sum polynomials p_1, p_2, \dots, p_m is same as the algebra generated by h_1, h_2, \dots, h_m , and also by e_1, e_2, \dots, e_m . Thus one has

$$\langle p_1, p_2, \dots, p_m \rangle = \langle h_1, h_2, \dots, h_m \rangle = \langle e_1, e_2, \dots, e_m \rangle. \quad (8)$$

One may also conclude (8) from Newton's formula (1) and (3). By [8, Theorem 4.3], $\langle p_1, p_2, \dots, p_m \rangle$ is a prime ideal in S . Thus, we can conclude that $\langle h_1, h_2, \dots, h_m \rangle$ and $\langle e_1, e_2, \dots, e_m \rangle$ are also prime ideals in S . \square

Remark 3.15. Let $n = 4$ in Proposition 3.14. Then h_1, h_2, h_a form a regular sequence in S provided $h_a \notin \langle h_1, h_2 \rangle$. By Newton's formula (1), we observe that $h_5 \equiv 0 \pmod{\langle h_1, h_2 \rangle}$. Hence $h_5 \in \langle h_1, h_2 \rangle$. We see that h_1, h_2 generate a prime ideal in S , but h_1, h_2, h_5 do not form a regular sequence in S . Thus, we need the condition $h_a \notin \langle h_1, h_2 \rangle$.

Computer calculations in CoCoA [2] suggest that $I = \langle h_1, h_{2m} \rangle$, where $m \in \mathbb{N}$, should be a prime ideal in $S = \mathbb{C}[x_1, x_2, x_3, x_4]$. For $m = 1$, it follows from Proposition 3.14. We prove for $m = 2$ in the following example.

Example 3.16. Let $S = \mathbb{C}[x_1, x_2, x_3, x_4]$ be a polynomial ring. Let $I = \langle h_1, h_4 \rangle$. Then I is a prime ideal in S .

Proof. Let $R = S/I$. We compute the Jacobian of I , say Jacobian is J . Let $J' = I_2(J)$, denotes the ideal generated by 2×2 minors of Jacobian. Also $\text{ht}(I) = 2$, since I is generated by a regular sequence of length 2. The determinants of 2×2 minors of the Jacobian can be written as

$$J' = \left\langle \frac{\partial h_4}{\partial x_i} - \frac{\partial h_4}{\partial x_j} \right\rangle \text{ for } 1 \leq i < j \leq 4.$$

By Lemma 2.1 (i), we may write J' as

$$J' = \langle (x_j - x_i)h_2 + (x_j^2 - x_i^2)h_1 + (x_j^3 - x_i^3) \rangle \text{ for } 1 \leq i < j \leq 4.$$

Thus, we have

$$I + J' = \langle h_1, h_4, (x_j - x_i)h_2 + (x_j^2 - x_i^2)h_1 + (x_j^3 - x_i^3) \rangle \text{ for } 1 \leq i < j \leq 4.$$

Claim: $\sqrt{I + J'} = (x_1, x_2, x_3, x_4)$.

Suppose not, that is, there exists $w = (w_1, w_2, w_3, w_4) \in \mathbb{P}^3$ with $w \in Z(I + J')$. We assume that none of w_i is zero. Also assume that $w_i \neq w_j$ for $i \neq j$. Since w is in \mathbb{P}^3 , we can make $w_1 = 1$ as $w_1 \neq 0$. So, let $w = (1, x, y, z)$. As $w \in Z(I + J')$, we have $h_1(w) = 0 = h_4(w)$, moreover

$$(x_j - x_i)h_2(w) + (x_j^3 - x_i^3) = 0 \text{ for } 1 \leq i < j \leq 4. \quad (9)$$

By (9), either $x_i = x_j$ or $h_2(w) = -(x_j^2 + x_i x_j + x_i^2)$ for $1 \leq i < j \leq 4$. By assumption $x_i \neq x_j$. Thus we have $h_2(w) = -(x_j^2 + x_i x_j + x_i^2)$ for $1 \leq i < j \leq 4$. An easy simplification shows that it is not possible. We may argue similarly when $w_i = w_j$ for $i \neq j$. Hence there is no nontrivial solution. By similar argument as used in Theorem 3.8, we conclude that R is a normal domain and I is a prime ideal. \square

Simply note that by [1, Lemma 2.2 and Proposition 2.9], the sequence of polynomials $p_a, p_{2a}, \dots, p_{na}$ and $h_a, h_{2a}, \dots, h_{na}$ form a regular sequence in $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ respectively. We partially extend the above conclusion in the following proposition:

Proposition 3.17. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring. Let $n, b, k \in \mathbb{N}$. Then the following holds:

- (i) $p_a, p_{2a}, \dots, p_{(n-1)a}, p_b$ form a regular sequence in S if and only if $b = nak$.
- (ii) $h_a, h_{2a}, \dots, h_{(n-1)a}, h_b$ form a regular sequences in S if and only if $b = nak$.

Proof. By Newton's formula (3), one has

$$p_c = \begin{cases} (-1)^k n e_n^k \pmod{(p_1, p_2, \dots, p_{n-1})}, & \text{if } c = nk; \\ 0 \pmod{(p_1, p_2, \dots, p_{n-1})}, & \text{otherwise.} \end{cases}$$

Clearly $p_1, p_2, \dots, p_{n-1}, p_c$ form a regular sequence in S if and only if $c = nk$. Thus the claim (i) follows from [1, Lemma 2.2]. By Newton's formula (1), one has

$$h_c = \begin{cases} (-1)^k e_n^k \pmod{(h_1, h_2, \dots, h_{n-1})}, & \text{if } c = nk; \\ 0 \pmod{(h_1, h_2, \dots, h_{n-1})}, & \text{otherwise.} \end{cases}$$

Clearly $h_1, h_2, \dots, h_{n-1}, h_c$ form a regular sequence in S if and only if $c = nk$. Thus the claim (ii) also follows from [1, Lemma 2.2]. \square

4. FINAL REMARKS

Following question is a special case of Question 1.1:

Question 4.1. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring with $n \geq 4$. Let $I = \langle p_a, p_b \rangle$, where $a, b \in \mathbb{N}$. For which pairs of integers a, b , I is a prime ideal in S .

We answer the previous question to some extent in the Theorem 3.8. We observe that the Theorem 3.8 is still in weaker form, due to arithmetic condition $q_1 > \max\{n, a\}$. In fact one can answer all the prime ideals which arises by using Serre criterion for normality and vanishing sums of roots of unity, by answering the following problem:

Problem 4.2. Let $m \in \mathbb{N}$. Consider m -th roots of unity in the field of complex number \mathbb{C} . For which natural numbers n and $k \geq 2$, do there exist m -th roots of unity $\alpha_1, \dots, \alpha_n \in \mathbb{C}$ such that $\alpha_1^k + \alpha_2^k + \dots + \alpha_n^k \neq 0$.

Similar to Question 4.1, one may also ask the following question:

Question 4.3. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring with $n \geq 4$. Let $I = \langle h_a, h_b \rangle$, where $a, b \in \mathbb{N}$. For which pairs of integers a, b , I is a prime ideal in S .

For $n = 4$ in the previous question, the computational calculations in CoCoA [2] suggest that $I = \langle h_1, h_{2m} \rangle$, where $m \in \mathbb{N}$, should be a prime ideal in $S = \mathbb{C}[x_1, x_2, x_3, x_4]$.

Recall the following definition:

Definition 4.4. Let $R = \bigoplus_{i=0}^c R_i$, $R_c \neq 0$ be a graded Artinian algebra. We say that R has the strong Lefschetz property (SLP) if there exists an element $L \in R_1$ such that the multiplication map

$$\times L^d : R_i \longrightarrow R_{i+1}$$

has full rank for all $0 \leq i \leq c-1$ and $1 \leq d \leq c-i$. We call an $L \in R_1$ with this property a strong Lefschetz element.

Let $J = \langle p_a, p_{a+1}, \dots, p_{a+n-1} \rangle$ in the polynomial ring $S = K[x_1, x_2, \dots, x_n]$ over a field K . Then the Artinian ring $R = S/J$ has the SLP, see [6, Proposition 7.1]. In the following example, we will show that for a complete intersection ideal $I = \langle h_a, h_{a+1}, \dots, h_{a+n-1} \rangle$, the Artinian ring $R = S/I$ have the SLP.

Example 4.5. Let K be a field of characteristic zero. Let $S = K[x_1, x_2, \dots, x_n]$ be a polynomial ring over K with standard grading, i.e. $\deg x_i = 1$ for all i . Let $I = \langle h_a, h_{a+1}, \dots, h_{a+n-1} \rangle$. Then $R = S/I$ has the SLP.

Proof. Consider the initial ideal of I with respect to lexicographic term order, one has

$$\text{in}(I) = \langle x_1^a, x_2^{a+1}, \dots, x_n^{a+n-1} \rangle.$$

Stanley [14] proved that every monomial complete intersection

$$K[x_1, x_2, \dots, x_n] / \langle x_1^a, x_2^{a+1}, \dots, x_n^{a+n-1} \rangle$$

has the SLP with $x_1 + x_2 + \dots + x_n$ as a strong Lefschetz element using the fact that it is isomorphic to the cohomology ring of a direct product of projective spaces over the complex number field. We see that $S/\text{in}(I)$ has the SLP. Thus, by [15, Proposition 2.9] we conclude that R has the SLP. \square

Question 4.6. Let $S = \mathbb{C}[x_1, x_2, \dots, x_n]$ be a polynomial ring. Let $k \in \mathbb{N}$. Is it true that for the ideal $I = \langle h_a, h_{2a}, \dots, h_{(n-1)a}, h_{nak} \rangle$, one has the initial ideal

$$\text{in}(I) = \langle x_1^a, x_2^{2a}, \dots, x_n^{nak} \rangle.$$

If the answer to the previous question is positive. Then again one can construct more examples of Artinian ring having the SLP. In a joint work with Martino [8], we explicitly derive several examples of complete intersection ideal generated by complete symmetric polynomials. Again, one can ask similar question for those complete intersection ideals.

We conclude the section with one remark from the recent paper of Fröberg and Shapiro [5], where the authors established a connection between regular sequences of complete symmetric polynomials and the codimension of the Vandermonde variety. To an arbitrary pair $(k; I)$, where $k \geq 2$ is a positive integer and $I = \{i_0 < i_1 < \dots < i_{m-1}\}$, $m \geq k$ is a sequence of integers, Fröberg and Shapiro discuss the Vandermonde variety $Vd_{k;I}^A$ in [5]. Under the assumption $i_0 = 0$ and $\gcd(i_1, \dots, i_{m-1}) = 1$, the authors asked [5, Problem 2], for which pairs $(k; I)$, the variety $Vd_{k;I}^A$ has the expected codimension. In the first non-trivial case $k = 3, m = 5$, the authors conclude that the variety $Vd_{3;I}^{21}$ has the expected codimension (equal to 3) if and only if three complete symmetric polynomials $h_{i_2-2}, h_{i_3-2}, h_{i_4-2}$ form a regular sequence in $\mathbb{C}[x_1, x_2, x_3]$. The problem of when three complete symmetric polynomials h_a, h_b, h_c , form a regular sequence in $\mathbb{C}[x_1, x_2, x_3]$ was considered in [1, Conjecture 2.17]. In [5, Conjecture 13], in a special case, it is mentioned that if $(a, b, c) = (1, 4, 3k + 2), k \geq 1$, then h_a, h_b, h_c neither is a regular sequence, nor $h_c \in (h_a, h_b)$. This will be clear from the following proposition:

Proposition 4.7. Let $S = \mathbb{C}[x_1, x_2, x_3]$ be a polynomial ring. Then h_1, h_4, h_n form a regular sequence in S if and only if $n = 3k, k \geq 1$.

Proof. By Newton's formula (1), one has

$$h_n = \begin{cases} e_3^k \pmod{(h_1, h_4)}, & \text{if } n = 3k; \\ 0 \pmod{(h_1, h_4)}, & \text{if } n = 3k + 1; \\ -(k+1)e_2e_3^k \pmod{(h_1, h_4)}, & \text{if } n = 3k + 2. \end{cases}$$

Thus the claim follows from [8, Theorem 2.2]. \square

REFERENCES

- [1] A. Conca, C. Krattenthaler, and J. Watanabe. *Regular sequences of symmetric polynomials*. Rend. Semin. Math. Univ. Padova **121**, 179–199 (2009).
- [2] The CoCoATeam. *CoCoA: a system for doing Computations in Commutative Algebra*. Available at <http://cocoa.dima.unige.it>.
- [3] R. Dvornicich and U. Zannier. *Newton functions generating symmetric fields and irreducibility of Schur polynomials*. Adv. Math. **222**, no. 6, 1982–2003 (2009).
- [4] D. Eisenbud. *Commutative Algebra*. With a view toward algebraic geometry. Graduate Texts in Mathematics, **150** Springer-Verlag, New York, xvi+785 pp. (1995).
- [5] R. Fröberg and B. Shapiro. *Vandermonde varieties and relations among Schur polynomials*. arXiv:1302.1298v1 [math.AC].
- [6] T. Harima and J. Watababe. *The strong Lefschetz property for Artinian algebras with non-standard grading*. J. Algebra. **311**, no. 2, 511–537 (2007).
- [7] M. Hochster. *Big Cohen-Macaulay modules and algebras and embeddability in rings of Witt vectors*. Queen’s Papers on Pure and Applied Math. **42**, 106–195 (1975).
- [8] N. Kumar and I. Martino. *Regular sequences of power sums and complete symmetric polynomials*. Matematiche (Catania). **67**, no. 1, 103–117 (2012).
- [9] T. Y. Lam and K.H. Leung. *On Vanishing sums of roots of unity*. Journal of algebra. **224**, no. 1, 91–109 (2000).
- [10] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Second edition. With contributions by A. Zelevinsky. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, x+475 pp, (1995).
- [11] H. B. Mann. *On linear relations between the roots of unity*. Mathematika. **12**, 107–117 (1965).
- [12] J. S. Milne. *Algebraic geometry*. Allied Publishers. (2005).
- [13] B. Poonen and M. Rubinstein. *The number of intersection points made by the diagonal of a regular polygon*. SIAM J. Discrete Math. **11**, no. 1, 135–156 (1998).
- [14] R. P. Stanley. *Weyl groups, the hard Lefschetz theorem, and the Sperner property*. SIAM J. Algebraic Discrete Methods. **1**, no. 2, 168–184 (1980).
- [15] A. Wiebe. *The Lefschetz property for componentwise linear ideals and Gotzmann ideals*. Comm. Algebra. **32**, no. 12, 4601–4611 (2004).

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI GENOVA, VIA DODECANESO 35, 16146 GENOVA, ITALY

E-mail address: kumar@dima.unige.it