

MA 109 D1&D2 Lecture 4

Ravi Raghunathan

Department of Mathematics

November 23, 2020

Cauchy sequences: the definition

The Prehistory of Limits

Cauchy sequences

As we have seen, it is not easy to tell whether a sequence converges or not because we have to first guess what the limit might be, and then try and prove that the sequence actually converges to this limit. For a monotonic sequence, we have a criterion, but what about more general sequences?

There is another very useful notion which allows us to decide whether the sequence converges **by looking only at the elements of the sequence itself**. We describe this below.

Definition: A sequence a_n in \mathbb{R} is said to be a **Cauchy sequence** if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon,$$

for all $m, n > N$.

Cauchy sequences: the theorem

Theorem 4: Every Cauchy sequence in \mathbb{R} converges (to a real number, obviously).

Remark 1: One can now check the convergence of a sequence just by looking at the sequence itself!

Remark 2: One can easily check the converse:

Theorem 5: Every convergent sequence (in any set X) is Cauchy.

Remark 3: Remember that when we defined sequences we defined them to be functions from \mathbb{N} to X , for any set X . So far we have only considered $X = \mathbb{R}$, but as we said earlier we can take other sets, for instance, subsets of \mathbb{R} . For instance, if we take $X = \mathbb{R} \setminus 0$, Theorem 4 is not valid. The sequence $1/n$ is a Cauchy sequence in this X but obviously does not converge in X . If we take $X = \mathbb{Q}$, the example given in 1.5.(i) ($a_{n+1} = (a_n + 2/a_n)/2$) is a Cauchy sequence in \mathbb{Q} which does not converge in \mathbb{Q} . Thus Theorem 4 is really a theorem about real numbers.

The completeness of \mathbb{R}

A set X in which every Cauchy sequence converges (to a limit in X) is called a complete set. Thus Theorem 4 is sometimes rewritten as

Theorem 4': The real numbers are complete.

We will see other examples of complete sets, but we can now address (very briefly) the question of what a real number is. More precisely, we can *construct* the set of real numbers starting with the rational numbers.

We let S be the set of all sequences with values in \mathbb{Q} . We will put a relation on this set.

The definition of a real number

Two sequence $\{a_n\}$ and $\{b_n\}$ will be related to each other (and we write $a_n \sim b_n$) if

$$\lim_{n \rightarrow \infty} |a_n - b_n| = 0$$

You can check that this is an equivalence relation and it is a fact that it *partitions* the set S into disjoint classes. The set of disjoint classes is denote S / \sim .

You can easily see that if two sequences converge to the same limit, they are necessarily in the same class.

Definition: A real number is an equivalence class in S / \sim .

So a real number should be thought of as the collection of all rational sequences which converge to it.

Sequences in \mathbb{R}^2 and \mathbb{R}^3

Most of our definitions for sequences in \mathbb{R} are actually valid for sequences in \mathbb{R}^2 and \mathbb{R}^3 . Indeed, the only thing we really need to define the limit is the notion of distance. Thus if we replace the modulus function $||$ on \mathbb{R} by the distance functions in \mathbb{R}^2 and \mathbb{R}^3 all the definitions of convergent sequences and Cauchy sequences remain the same.

For instance, a sequence $a(n) = (a(n)_1, a(n)_2)$ in \mathbb{R}^2 is said to converge to a point $l = (l_1, l_2)$ (in \mathbb{R}^2) if for all $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$\sqrt{(a(n)_1 - l_1)^2 + (a(n)_2 - l_2)^2} < \epsilon$$

whenever $n > N$. A similar definition can be made in \mathbb{R}^3 using the distance function on \mathbb{R}^3 .

Theorems 2 (the Sandwich Theorem) and 3 (about monotonic sequences) don't really make sense for \mathbb{R}^2 or \mathbb{R}^3 because there is no ordering on these sets, that is, it doesn't really make sense to ask if one point on the plane or in space is less than the other.



The first man to think about limits?



Zeno of Elea (490 - 460 BCE) was a famous Greek philosopher (source: Wikipedia)

Zeno of Elea

First let us record that we have no idea what Zeno looked like. The picture above was painted in the period 1588 - 1594 CE in Spain, about two thousand years after Zeno's time. Here are two more images of Zeno (also from Wikipedia)



Zeno's Paradoxes

I couldn't find out where the first statue came from and when it was made. The second seems to have come from Herculaneum in Italy (incidentally, Elea (modern Vilia) is a town in Italy). Now Herculaneum was destroyed by a volcanic eruption from the nearby volcano Vesuvius in 79 CE, so it looks like the bust was created within 500 years of Zeno's death. Maybe it was even made during his lifetime and was lying around in some wealthy Roman's house for the next few centuries. Unfortunately, it is not clear whether this statue is one of Zeno of Elea or of another Zeno (of Citium) who lived about 150 years later. So we still really have no clue how he looked.

The important about Zeno is that it would appear that he was the first human to think about limits and limiting processes, at least in recorded history. Most of what we know about him is through his paradoxes, nine of which survive in the works of another famous Greek philosopher Aristotle (384 - 322 CE) , the official guru/tutor of Alexander the Great (aka Sikander in India).

Achilles and the tortoise

One of Zeno's motivations for stating his paradoxes seems to have been to defend his own guru Parmenides' philosophy (whatever that was). Anyway here is his most famous paradox as recorded by Aristotle.

Achilles and the tortoise:

In a race, the quickest runner can never overtake the slowest, since the pursuer must first reach the point whence the pursued started, so that the slower must always hold a lead.

Aristotle, Physics VI:9, 239b15

General knowledge question: Who was Achilles?

Zeno's paradox animated

Achilles and the Tortoise

A gateway to infinite series

Nowadays, this line of argument does not really bother us, since we understand that an infinite number of terms (in this case consisting of the time travelled in each segment or the distance travelled in each segment) can add up to something finite.

Nevertheless there are other philosophical issues that continued to bother mathematicians and physicists for a long time. After all, this kind of discussion does lead us to question whether intervals of time and space can be infinitely subdivided, or if “instantaneous motion” makes sense.

Since we are learning mathematics, we won't speculate on physics or philosophy, but we note that Zeno's argument gives a good way to derive the sum of an infinite geometric series. The geometric series is one of the simplest examples of infinite series, so let us see how this is done.

Geometric series - the formula

Let us suppose that the speed of achilles is v and that the speed of the tortoise is rv for some $0 < r < 1$. We will assume that the tortoise was given a headstart of distance “ a ”.

- ▶ The distance covered by Achilles in time t is vt .
- ▶ The distance covered by the tortoise in time t is rvt .
- ▶ Achilles catches up with the tortoise when $vt = a + rvt$, that is, at time $t = a/(v - rv)$ and when the total distance covered by Achilles is $vt = a/(1 - r)$.

On the other hand,

- ▶ Distance covered by the tortoise by the time Achilles has covered distance a is ar .
- ▶ Distance covered by the tortoise by the time Achilles has covered distance ar is ar^2
- ▶ Total distance covered by Achilles when he has caught up with the tortoise is $a + ar + ar^2 + \dots$.
- ▶ Thus we get $a + ar + ar^2 + \dots = a/(1 - r)$.

Infinite series - a more rigorous treatment

Let us recall what we mean when we write

$$a + ar + ar^2 + \dots = \frac{a}{1-r}.$$

Another way of writing the same expression is

$$\sum_{k=0}^{\infty} ar^k = \frac{a}{1-r}.$$

The precise meaning is the following. Form the **partial sums**

$$s_n = \sum_{k=0}^n ar^k.$$

These partial sums $s_1, s_2, \dots, s_n, \dots$ form a sequence and by

$\sum_{k=0}^{\infty} ar^k = a/(1-r)$, we mean $\lim_{n \rightarrow \infty} s_n = a/1-r$.

So when we speak of the sum of an infinite series, what we really mean is the limit of its partial sums.

Convergence of the geometric series

So to justify our formula we should show that $\lim_{n \rightarrow \infty} s_n = a/(1-r)$, that is, given $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$\left| s_n - \frac{a}{1-r} \right| < \epsilon,$$

for all $n > N$.

In other words we need to show that

$$\left| \frac{a(1-r^{n+1})}{1-r} - \frac{a}{1-r} \right| = \left| \frac{ar^{n+1}}{1-r} \right| < \epsilon$$

if n is chosen large enough.

But $\lim_{n \rightarrow \infty} r^n = 0$, so there exists N such that $r^{n+1} < (1-r)\epsilon/a$ for all $n > N$, so for this N , if $n > N$,

$$\left| s_n - \frac{a}{1-r} \right| < \epsilon.$$

This shows that the geometric series converges to the given expression.