

MA 105 D1 Lecture 4

Ravi Raghunathan

Department of Mathematics

August 3, 2017

Recap

More about continuous functions

Arnold's problem

Functions of several variables

Set Cardinality

Continuity - the definition

Definition: If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be **continuous at the point c** if and only if

$$\lim_{x \rightarrow c} f(x) = f(c).$$

Thus, if c is one of the end points we require only the left or right hand limit to exist.

A function f on (a, b) (resp. $[a, b]$) is said to be **continuous** if and only if it is continuous at every point c in (a, b) (resp. $[a, b]$).

If f is not continuous at a point c we say that it is **discontinuous at c** , or that **c is a point of discontinuity for f** .

Intuitively, continuous functions are functions whose graphs can be drawn on a sheet of paper without lifting the pencil of the sheet of paper. That is, there should be no “jumps” in the graph of the function.

The composition of continuous functions

Theorem 8: Let $f : (a, b) \rightarrow (c, d)$ and $g : (c, d) \rightarrow (e, f)$ be functions such that f is continuous at x_0 in (a, b) and g is continuous at $f(x_0) = y_0$ in (c, d) . Then the function $g(f(x))$ (also written as $g \circ f(x)$ sometimes) is continuous at x_0 . So the composition of continuous functions is continuous.

Exercise 4: Prove the theorem above starting from the definition of continuity.

Using the theorem above we can show that $\cos x$ is continuous if we show that \sqrt{x} is continuous, since $\cos x = \sqrt{1 - \sin^2 x}$ and we know that $1 - \sin^2 x$ is continuous since it is the product of the sums of two continuous functions ($(1 + \sin x)$ and $(1 - \sin x)$!).

Once we have the continuity of $\cos x$ we get the continuity of all the rational trigonometric functions, that is functions of the form $P(x)/Q(x)$, where P and Q are polynomials in $\sin x$ and $\cos x$, provided $Q(x)$ is not zero.

The continuity of the square root function

Thus in order to prove the continuity of $\cos x$ (assuming the continuity of $\sin x$) we need only prove the continuity of the square root function.

The main observation is that continuity is a **local** property, that is, **only the behaviour of the function near the point being investigated is important**.

Let $x_0 \in [0, \infty)$. To show that the square root function is continuous at x_0 we need to show that $\lim_{y \rightarrow x_0} \sqrt{y} = \sqrt{x_0}$, that is we need to show that $|\sqrt{y} - \sqrt{x_0}| < \epsilon$ whenever $0 < |y - x_0| < \delta$ for some δ . First assume that $x_0 \neq 0$. Then

$$|\sqrt{y} - \sqrt{x_0}| = \left| \frac{y - x_0}{\sqrt{y} + \sqrt{x_0}} \right| < \frac{|y - x_0|}{\sqrt{x_0}}.$$

If we choose $\delta = \epsilon\sqrt{x_0}$, we see that

$$|\sqrt{y} - \sqrt{x_0}| < \epsilon,$$

which is what we needed to prove. When $x_0 = 0$, I leave the proof as an exercise. □

The intermediate value theorem

One of the most important properties of continuous functions is the Intermediate Value Property (IVP). We will use this property repeatedly to prove other results.

Theorem 9: Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. For every u between $f(a)$ and $f(b)$ there exists $c \in [a, b]$ such that $f(c) = u$.

Functions which have this property are said to have the Intermediate Value Property. Theorem 9 can thus be restated as saying that continuous functions have the IVP.

We will not be proving this property - it is a consequence of the completeness of the real numbers. Intuitively, this is clear. Since one can draw the graph of the function without lifting one's pencil off the sheet of paper, the pencil must cut every line $y = e$ with e between $f(a)$ and $f(b)$.

The IVT in a picture




Image created by Enoch Lau see

<http://en.wikipedia.org/wiki/File:Intermediatevaluethorem.png>

(Creative Commons Attribution-Share Alike 3.0 Unported license).

Zeros of functions

One of the most useful applications of the intermediate value property is to find roots of polynomials, or, more generally, to find zeros of continuous functions, that is to find points $x \in \mathbb{R}$ such that $f(x) = 0$.

Theorem 10: Every polynomial of odd degree has at least one real root.

Proof: Let $P(x) = a_n x^n + \dots + a_0$ be a polynomial of odd degree. We can assume without loss of generality that $a_n > 0$. It is easy to see that if we take $x = b > 0$ large enough, $P(b)$ will be positive. On the other hand, by taking $x = a < 0$ small enough, we can ensure that $P(a) < 0$. Since $P(x)$ is continuous, it has the IVP, so there must be a point $x_0 \in (a, b)$ such that $f(x_0) = 0$. \square

The IVP can often be used to get more specific information. For instance, it is not hard to see that the polynomial $x^4 - 2x^3 + x^2 + x - 3$ has a root that lies between 1 and 2.

Continuous functions on closed, bounded intervals

The other major result on continuous functions that we need is the following. A closed bounded interval is one of the form $[a, b]$, where $-\infty < a$ and $b < \infty$.

Theorem 11: A continuous function on a closed bounded interval $[a, b]$ is bounded and attains its infimum and supremum, that is, there are points x_1 and x_2 in $[a, b]$ such that $f(x_1) = m$ and $f(x_2) = M$, where m and M denote the infimum and supremum respectively.

Again, we will not prove this, but will use it quite often. Note the contrast with open intervals. The function $1/x$ on $(0, 1)$ does not attain a maximum - in fact it is unbounded. Similarly the function $1/x$ on $(1, \infty)$ does not attain its minimum, although, it is bounded below.

Exercise 5: In light of the above theorem, can you find a **continuous** function $g : (a, b) \rightarrow \mathbb{R}$ for part (i) of Exercise 1.11, with $c \in (a, b)$?

The function $\sin \frac{1}{x}$

Let us look at Exercise 1.13 part (i).

Consider the function defined as $f(x) = \sin \frac{1}{x}$ when $x \neq 0$, and $f(0) = 0$. The question asks if this function is continuous at $x = 0$. How about $x \neq 0$? Why is $f(x)$ continuous? Because it is a composition of the sin function and a rational function $1/x$. Since both of these are continuous when $x \neq 0$, so is $f(x)$.

Let us look at the sequence of points $x_n = 2/(2n + 1)\pi$. Clearly $x_n \rightarrow 0$ as $n \rightarrow \infty$.

For these points $f(x_n) = \pm 1$. This means that no matter how small I take my δ , there will be a point $x_n \in (0, \delta)$, such that $|f(x_n)| = 1$. But this means that $|f(x) - f(0)| = |f(x)|$ cannot be made smaller than 1 no matter how small δ may be. Hence, f is not continuous at 0. The same kind of argument will show that there is no value that we can assign $f(0)$ to make the function $f(x)$ continuous at 0.

You can easily check that $f(x)$ has the IVP. However, we have proved that it is not continuous. So IVP \nrightarrow continuity.

Sequential continuity

The preceding example showed that in order to demonstrate that a function, say $f(x)$, is not continuous at a point x_0 it is enough to find a sequence x_n tending to x_0 such that the value of the function $|f(x_n) - f(x_0)|$ remains large. Suppose it is not possible to find such a sequence. Does that mean the function is continuous at x_0 ? The following theorem answers the question affirmatively.

Theorem 12: A function $f(x)$ is continuous at a point a if and only if **for every sequence $x_n \rightarrow a$** , $\lim_{x_n \rightarrow a} f(x_n) = f(a)$.

A function that satisfies the property that for every sequence $x_n \rightarrow a$, $\lim_{x_n \rightarrow a} f(x_n) = f(a)$ is said to be **sequentially continuous**. The theorem says that sequential continuity and continuity are the same thing. Indeed, it is clear that a continuous function is necessarily sequentially continuous. It is the reverse that is slightly harder to prove.

Limits of functions of several variables

Just like we did for sequences in \mathbb{R}^2 and \mathbb{R}^3 , we can define the notion of the limit of a function for functions from \mathbb{R}^2 to \mathbb{R} .

Let $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ be a function. The function $f(x_1, x_2)$ is said to tend to a limit l as $(x_1, x_2) \rightarrow (a_1, a_2)$ if for any $\epsilon > 0$ there exists $\delta > 0$ such that

$$|f(x_1, x_2) - l| < \epsilon$$

whenever $0 < \sqrt{(x_1 - a_1)^2 + (x_2 - a_2)^2} < \delta$. Notice, that one can now approach the point (a_1, a_2) from any direction in the plane. Our definition requires that the limits from the different directions all exist and be equal. This is quite a powerful condition.

If we have functions from $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ we can make exactly the same definition. But this time $l = (l_1, l_2)$ will be in \mathbb{R}^2 and so will $f(x_1, x_2) = (f_1(x_1, x_2), f_2(x_1, x_2))$, so we will have to replace the modulus function by the distance between these two quantities:

$$\sqrt{[f_1(x_1, x_2) - l_1]^2 + [f_2(x_1, x_2) - l_2]^2}.$$

Limits of functions of several variables

The definitions we have made go through for functions from \mathbb{R}^m to \mathbb{R}^n , where m and n may be different. For instance, we have considered the case when $m = 2$ and $n = 1$ and also the case $m = 2$ and $n = 2$ above. But we could allow m and n to take any of the values 1, 2 or 3 (in fact, we can allow values greater than 3 as well!).

Exercise 1: Show that

$$\lim_{y \rightarrow x} f(y) = l$$

for $f = (f_1, f_2) : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \iff$

$$\lim_{y \rightarrow x} f_1(y) = l_1 \quad \text{and} \quad \lim_{y \rightarrow x} f_2(y) = l_2,$$

where $l = (l_1, l_2)$. In other words, when dealing with limits of functions which are vector-valued, it is enough to study the limits of the coordinate functions.

Continuous functions of several variables

Once the definition of the limit is clear it makes sense to talk of continuity as well. All the definitions remain the same, only the definition of the distance function changes depending on the domain and the range.

For instance, provided we know what “closed and bounded sets” are in \mathbb{R}^2 or \mathbb{R}^3 , Theorem 11 goes through for continuous functions $f : \mathbb{R}^m \rightarrow \mathbb{R}$. ($m = 2, 3$). For functions with more than one variable in the range the first part of Theorem 11 still works, but for the second part things are more complicated (again there is no “ordering” in \mathbb{R}^2 or \mathbb{R}^3).

While it is easy to see what a bounded set in \mathbb{R}^m should be, closed is a little more complicated and we will not give the definition here. However, a rectangle of the form $[a, b] \times [c, d]$ in \mathbb{R}^2 is an example of a closed and bounded set (also called “compact sets” of this form).

Theorem 12 goes through without any problems even when the range is in \mathbb{R}^2 or \mathbb{R}^3 .

Counting versus pairing

Having names for and dealing with large numbers (lakhs, millions, crores, billions etc.) is a very recent phenomenon. For most of human history, people did not have names for numbers beyond one hundred or a thousand. Indeed, supposedly some “primitive” tribes only had names for “one” and “two”. Everything else was “many”.

Nevertheless if you gave a stone-age man a herd of cattle and flock of sheep and asked him which had more animals, he could certainly have told you even if he didn't have names for the numbers of the cattle and the sheep. How?

He would simply have taken one cow from the herd and paired it with one sheep from the flock, and he would have continued to pair the animals in twos until either the cattle or the sheep ran out.

What do we mean by a cardinal number?

If the cows and sheep are exactly paired our stone-age man can say that there are the same number of cows and sheep.

Note that he does not need to count the number of cows individually (or count the number of sheep). He does not need names for numbers at all!

So what do we mean by a cardinal number? For instance, what do we mean by “seven”. It is the name we give for a certain property of various collections of objects- cows, sheep, books, ships, colours, chemical compounds, grains of rice. The objects of any one collection can be paired off exactly with objects in one of the other collections.

Cardinality: the definition

So “seven” is the property that all possible collections with a certain number of objects (namely, seven!) have. It is a completely abstract concept. “Sevenness” is the property that all of these sets of objects have.

Definition: Two sets X and Y are said to have the same cardinality if there is a bijective function from X to Y .

All of this discussion probably feels obvious to you. The discussion gets more tricky (and thus more interesting!) when X and Y are infinite sets.

Infinite sets

Example: Let $X = \mathbb{N}$ and let Y be set of all perfect squares of numbers in \mathbb{N} . Clearly X and Y have the same cardinality since the function $f(n) = n^2$ gives a bijective map from X to Y .

But Y is a subset of X ! So a set and a proper subset can have the same cardinality! This could never happen if X and Y were finite!

A set that has the same cardinality as the set \mathbb{N} is called **countable** or **denumerable**. Thus the set of perfect squares is countable. How about the set of odd natural numbers? The set of integers? The set of rational numbers \mathbb{Q} ?