

**MA 2017, Tutorial Sheet-5**  
**Boundary value problem and Fourier expansion**

1. Show that  $\sum_{n=1}^{\infty} \frac{1}{n} \sin nx \sin^2 n\alpha = \begin{cases} \text{constant} & (0 < x < 2\alpha) \\ 0 & (2\alpha < x < \pi) \end{cases}$
2. Show that  $\sum_{n=1}^{\infty} \frac{(-1)^{n-1} \cos nx}{n^2} = \frac{\pi^2}{12} - \frac{x^2}{4}, \quad (-\pi \leq x \leq \pi).$
3. Show that  $\sum_{n=0}^{\infty} \frac{\sin(2n+1)x}{(2n+1)^3} = \frac{1}{8}\pi x(\pi - x), \quad (0 \leq x \leq \pi).$
4. Use the Fourier expansions given in problems (1), (2) and (3) along with Fourier's Theorem to deduce the following results.
  - (a)  $1 + \frac{1}{2} - \frac{1}{4} - \frac{1}{5} + \frac{1}{7} + \frac{1}{8} - \frac{1}{10} - \frac{1}{11} + \dots = \frac{2\pi}{3\sqrt{3}}$
  - (b)  $1 - \frac{1}{2} + \frac{1}{4} - \frac{1}{5} + \frac{1}{7} - \frac{1}{8} + \frac{1}{10} - \frac{1}{11} + \dots = \frac{\pi}{3\sqrt{3}}$
  - (c)  $1 - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \frac{1}{5^2} - \frac{1}{6^2} + \dots = \frac{\pi^2}{12}$
  - (d)  $1 + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \frac{1}{5^2} + \frac{1}{6^2} + \dots = \frac{\pi^2}{6}$  (Euler's formula)
  - (e)  $1 - \frac{1}{3^3} + \frac{1}{5^3} - \frac{1}{7^3} + \frac{1}{9^3} - \dots = \frac{\pi^3}{32}$
  - (f)  $1 + \frac{1}{3^2} + \frac{1}{5^2} + \frac{1}{7^2} + \frac{1}{9^2} + \dots = \frac{\pi^2}{8}$
  - (g)  $\frac{1}{1.3} - \frac{1}{3.5} + \frac{1}{5.7} - \frac{1}{7.9} + \dots = \frac{\pi}{4} - \frac{1}{2}$

5. Find the Fourier series of  $f(x)$  on  $[-L, L]$  and determine the value that the series takes for  $-L \leq x \leq L$ .

(a)  $L = \pi, f(x) = 2x - 3x^2,$       (b)  $L = 1, f(x) = 1 - 3x^2,$

(c)  $L = \pi, f(x) = |\sin x|,$

(f)  $L = 1, f(x) = \begin{cases} 0, & -1 < x < -1/2, \quad 1/2 < x < 1, \\ \cos \pi x, & -1/2 < x < 1/2, \end{cases}$ ,

(g)  $L = \pi,$  and  $f(x)$  is one of the following functions (i)  $e^x,$  (ii)  $(x - \pi) \cos x,$  (iii)  $\sin kx,$   $k$  not an integer.

(h)  $L = \pi, f(x) = x + |x|$

(i)  $L = \pi, f(x) = \begin{cases} -x & -\pi \leq x < 0 \\ x & 0 \leq x < \pi \end{cases}$

(j)  $L = 1, f(x) = \begin{cases} 0 & -1 < x < 0, \\ x & 0 < x < 1. \end{cases}$

(k)  $L = 1, f(x) = \begin{cases} -1, & -1 \leq x < 0 \\ 1, & 0 \leq x \leq 1 \end{cases}$

6. Expand each of the following functions in a Fourier cosine series on  $[0, L]$ .

(a)  $L = 1$ ,  $f(x) = e^{-x}$ ,

(b)  $L = 2$ ,  $f(x) = \begin{cases} 0, & 0 \leq x \leq 1 \\ 1, & 1 \leq x \leq 2 \end{cases}$ ,

(c)  $L = \pi$ ,  $f(x) = 2 \sin x \cos x$ ,

(d)  $f(x) = x^2 - L^2$ .

(e)  $f(x) = 3x^2(x^2 - 2L^2)$ ,

(f)  $f(x) = x^3(3x - 4L)$ ,

(g)  $x^2(3x^2 - 8Lx + 6L^2)$ .

7. Expand each of the following functions in a Fourier sine series on  $[0, L]$ .

(a)  $L = 1$ ,  $f(x) = e^{-x}$ ,

(b)  $L = 2a$ ,  $f(x) = \begin{cases} x, & 0 < x < a \\ a, & a \leq x \leq 2a \end{cases}$ ,

(c)  $L = \pi$   $f(x) = 2 \sin x \cos x$ ,

(d)  $L = \pi$   $f(x) = \cos x$ .

(e)  $f(x) = x(L^2 - x^2)$

(f)  $f(x) = x(x^3 - 2Lx^2 + L^3)$ ,

(g)  $f(x) = x(3x^4 - 5Lx^3 + 2L^4)$ .