MA108 ODE: Linear DE-s of Second Order

Preeti Raman
IIT Bombay

Non-homogeneous Second Order Linear ODE's

Consider the non-homogeneous DE

$$y'' + p(t)y' + q(t)y = r(t)$$

where $p(t), q(t), r(t)$ are continuous functions on an interval I. The associated homogeneous DE is

$$y'' + p(t)y' + q(t)y = 0.$$

Can we relate the solutions of the above two DE's?

Theorem

Let $f(t)$ be any solution of

$$y'' + p(t)y' + q(t)y = r(t)$$

and $y_1(t), y_2(t)$ be a basis of the solution space of the corresponding homogeneous DE. Then the set of solutions of the non-homogeneous DE is

$$\{c_1y_1(t) + c_2y_2(t) + f(t) \mid c_1, c_2 \in \mathbb{R}\}.$$

Proof: Let $\phi(t)$ be any solution of

$$L(y) = y'' + p(t)y' + q(t)y = r(t).$$

Then,

$$L(\phi(t) - f(t)) = L(\phi(t)) - L(f(t)) = r(t) - r(t) = 0.$$

Hence, $\phi(t) - f(t)$ is a solution of the homogeneous DE. Thus,

$$\phi(t) - f(t) = c_1 y_1(t) + c_2 y_2(t),$$

for $c_1, c_2 \in \mathbb{R}$. Hence,

$$\phi(t) = c_1 y_1(t) + c_2 y_2(t) + f(t).$$

Summary: In order to find the general solution of a non-homogeneous DE, we need to

- get one particular solution of the non-homogeneous DE
- get the general solution of the corresponding homogeneous DE.

Method of Undetermined Coefficients

Suppose the non-homogeneous ODE has constant coefficients. In this case, we know how to write down the general solution of the corresponding homogeneous ODE. So we need to find one solution of the non-homogeneous DE. One way to do this is called the method of undetermined coefficients. Thus, we have:

$$y'' + py' + qy = r(t),$$

with $p, q \in \mathbb{R}$, and r is a continuous function on I. The method of undetermined coefficients works if $r(t)$ involves $e^{at}, \sin at, \cos at$ or polynomials in t.

Method of Undetermined Coefficients

Example: Find a particular solution of the DE:

$$y'' - 3y' - 4y = 3e^{2t}.$$

We'll search for a solution of the form ae^{2t}, where a is a constant. So put $y = ae^{2t}$. We get:

$$(ae^{2t})'' - 3(ae^{2t})' - 4ae^{2t} = 3e^{2t}.$$

Thus,

$$4ae^{2t} - 6ae^{2t} - 4ae^{2t} = 3e^{2t}.$$

Thus,

$$a = -\frac{1}{2}.$$

Hence $-\frac{1}{2}e^{2t}$ is a particular solution of the DE.
How do you get the general solution? Analyse roots of $m^2 - 3m - 4 = 0$. So general solution is

$$y = c_1 e^{4t} + c_2 e^{-t} - \frac{1}{2}e^{2t}.$$

Method of Undetermined Coefficients

Consider the DE

$$y'' + py' + qy = r(t).$$

where p and q are real numbers. If

$$r(t) = r_1(t) + r_2(t) + \ldots + r_n(t),$$

where $r_i(t)$ are e^{at} or $\sin at$ or $\cos at$ or polynomials in t, consider the n subproblems

$$y'' + py' + qy = r_i(t), 1 \leq i \leq n.$$

If $y_i(t)$ is a particular solution of this problem, then,

$$y(t) = y_1(t) + y_2(t) + \ldots + y_n(t)$$

is a particular solution of

$$y'' + py' + qy = r(t).$$

Method of Undetermined Coefficients

Example: Find a particular solution of

$$y'' - 3y' - 4y = 2\sin t.$$

Make a guess as to functions of which form we'll search for as a solution. $a\sin t$? No. $a\sin t + b\cos t$? Yes. So set

$$y(t) = a\sin t + b\cos t.$$

Thus,

$$y' = a\cos t - b\sin t;\ \ y'' = -a\sin t - b\cos t.$$

Substituting, we get:

$$(-5a + 3b - 2)\sin t + (-3a - 5b)\cos t = 0.$$

Thus,

$$-5a + 3b = 2;\ \ 3a + 5b = 0$$

(Why?). Thus, $a = -\frac{5}{17},\ \ b = \frac{3}{17}$, and a particular solution is

$$y(t) = -\frac{5}{17}\sin t + \frac{3}{17}\cos t.$$

Method of Undetermined Coefficients

Example: Find a particular solution of

$$y'' - 3y' - 4y = 4t^2 - 1.$$

Set

$$y(t) = at^2 + bt + c.$$

Substituting, we get:

$$2a - 3(2at + b) - 4(at^2 + bt + c) = 4t^2 - 1,$$

i.e.,

$$-4at^2 + (-6a - 4b)t + (2a - 3b - 4c) = 4t^2 - 1.$$

Thus,

$$-4a = 4, \quad -6a - 4b = 0, \quad 2a - 3b - 4c = -1.$$

Thus,

$$a = -1, \quad b = \frac{3}{2}, \quad c = -\frac{11}{8}.$$

Thus, a particular solution is

$$y(t) = -t^2 + \frac{3}{2}t - \frac{11}{8}.$$

Method of Undetermined Coefficients

Example: Find a particular solution of

$$y'' - 3y' - 4y = -8e^t \cos 2t.$$

We should search for a solution of the form

$$y(t) = ae^t \cos 2t + be^t \sin 2t.$$

Then,

$$y'(t) = (a + 2b)e^t \cos 2t + (-2a + b)e^t \sin 2t,$$

and

$$y'' = (-3a + 4b)e^t \cos 2t + (-4a - 3b)e^t \sin 2t.$$

Substituting, we get:

$$-10a - 2b = -8, \ 2a - 10b = 0.$$

It follows that

$$a = \frac{10}{13} \text{ and } b = \frac{2}{13}.$$

Thus, a particular solution is

$$y(t) = \frac{10}{13}e^t \cos 2t + \frac{2}{13}e^t \sin 2t.$$

Method of Undetermined Coefficients

Example: Find a particular solution of

$$y'' + 4y = 3\cos 2t.$$

Since $r(t) = 3\cos 2t$, you would look for solutions of the form

$$y(t) = a\cos 2t + b\sin 2t.$$

Thus,

$$y'(t) = -2a\sin 2t + 2b\cos 2t,$$

$$y''(t) = -4a\cos 2t - 4b\sin 2t.$$

Substituting in the given DE, we get:

$$(-4a\cos 2t - 4b\sin 2t) + 4(a\cos 2t + b\sin 2t) = 3\cos 2t.$$

But the lhs is 0! So can't solve for a and b.

Method of Undetermined Coefficients

Why this ...? The problem was that $\sin 2t$ and $\cos 2t$ are also solutions of the associated homogeneous ODE: $y'' + 4y = 0$. So lesson learnt? When we search for solutions of a particular form, we need to make sure that it's not a solution of the associated homogeneous equation.
We now modify the proposed solution as:

$$y(t) = at\cos 2t + bt\sin 2t.$$

Then,

$$y'(t) = (b - 2at)\sin 2t + (a + 2bt)\cos 2t,$$

$$y''(t) = -4at\cos 2t - 4bt\sin 2t - 4a\sin 2t + 4b\cos 2t.$$

Substituting, we get:

$$-4a\sin 2t + 4b\cos 2t = 3\cos 2t.$$

Thus, $a = 0,\ b = \frac{3}{4}$, and a particular solution is $y(t) = \frac{3}{4}t\sin 2t$.

Method of Undetermined Coefficients

If the obvious candidate for a solution, say $y(t) = f(t)$, as well as this one multiplied by t, $y(t) = tf(t)$, turn out to be solutions of the associated homogeneous ODE, then what to do? Modify the proposed solution by multiplying it with t^2; i.e., set

$$y(t) = t^2 f(t).$$

Can this too be a solution of the homogeneous ODE? No, since the solution space is two dimensional.

Method of Undetermined Coefficients

Example: Find a particular solution of

$$y'' - 3y' - 4y = 3e^{2t} + 2\sin t + 4t^2 - 1 - 8e^t \cos 2t.$$

Here,

$$r(t) = r_1(t) + r_2(t) + r_3(t) + r_4(t).$$

We need to solve

$$y'' - 3y' - 4y = r_i(t),$$

get a particular solution $y_i(t)$, and then

$$y(t) = y_1(t) + y_2(t) + y_3(t) + y_4(t)$$

is a particular solution of the given problem. Thus, a particular solution is

$$y(t) = -\frac{1}{2}e^{2t} - \frac{5}{17}\sin t + \frac{3}{17}\cos t - t^2 + \frac{3}{2}t - \frac{11}{8} + \frac{10}{13}e^t \cos 2t + \frac{2}{13}e^t \sin 2t.$$

Method of Undetermined Coefficients

Example: Find a particular solution of

$$y'' + y = x^3 \sin x.$$

What's your candidate? Presence of $\sin x$ indicates both $\sin x$ and $\cos x$ in the answer. Presence of x^3 indicates a generic cubic polynomial. Thus,

$$(a_1x^3 + b_1x^2 + c_1x + d_1)\cos x + (a_2x^3 + b_2x^2 + c_2x + d_2)\sin x?$$

This wouldn't do since $\sin x$ and $\cos x$ are already solutions of the homogeneous part. So work with

$$x(a_1x^3 + b_1x^2 + c_1x + d_1)\cos x + x(a_2x^3 + b_2x^2 + c_2x + d_2)\sin x.$$

Method of Undetermined Coefficients

Consider the DE

$$y'' + py' + qy = r(t).$$

where p and q are real numbers.

Step I: Find the general solution of the homogeneous equation

$$y'' + py' + qy = 0.$$

Step II: If $r(t) = r_1(t) + r_2(t) + \ldots + r_n(t)$, where $r_i(t)$ are e^{at} or $\sin at$ or $\cos at$ or polynomials in t, consider

$$y'' + py' + qy = r_i(t), 1 \leq i \leq n.$$

Step III: Find a particular solution $y_i(t)$ for each of the above n subproblems. Then

$$y(t) = y_1(t) + y_2(t) + \ldots + y_n(t)$$

is a particular solution of

$$y'' + py' + qy = r(t).$$

Step IV: In case, a particular solution is same as the solution of the associated homogeneous equation, then we modify that particular solution by multiplying it with t or atmost t^2.

Method of Undetermined Coefficients

Table of particular solutions

$r(t)$	Particular solutions $y(t)$	
$P_n(t) = \sum_{i=0}^{n} a_i t^i$	$t^s \left(\sum_{i=0}^{n} A_i t^i\right)$	
$P_n(t)e^{\alpha t}$	$e^{\alpha t} t^s \left(\sum_{i=0}^{n} A_i t^i\right)$	
$P_n(t)e^{\alpha t}\sin\beta t$	$e^{\alpha t} t^s \left(\sum_{i=0}^{n} A_i t^i\right)\cos\beta t$ $e^{\alpha t} t^s \left(\sum_{i=0}^{n} B_i t^i\right)\sin\beta t$	$+$
$P_n(t)e^{\alpha t}\cos\beta t$	$e^{\alpha t} t^s \left(\sum_{i=0}^{n} A_i t^i\right)\cos\beta t$ $e^{\alpha t} t^s \left(\sum_{i=0}^{n} B_i t^i\right)\sin\beta t$	$+$

Here $s = 0, 1$ or 2 is the smallest integer so that no term in $y(t)$ is a solution of the associated homogeneous equation.

Method of Variation of Parameters

Consider the DE

$$y'' + p(x)y' + q(x)y = r(x),$$

where p, q and r are continuous on an interval I. The associated homogeneous DE is

$$y'' + p(x)y' + q(x)y = 0. \tag{1}$$

Suppose that we know the general solution

$$y(x) = c_1 y_1(x) + c_2 y_2(x)$$

of equation (1). In the method of variation of parameters, we replace the constants c_1, c_2 by functions $v_1(x), v_2(x)$, (to be suitably determined) so that

$$y(x) = v_1(x)y_1(x) + v_2(x)y_2(x)$$

is a solution of

$$y'' + p(x)y' + q(x)y = r(x).$$

$$y(x) = v_1(x)y_1(x) + v_2(x)y_2(x)$$

$$y' = v_1y_1' + v_2y_2' + v_1'y_1 + v_2'y_2.$$

Even though $v_1(x)$ and $v_2(x)$ are non-constant functions we would like them to behave like constants during product differentiation, i.e., we demand that

$$v_1'y_1 + v_2'y_2 = 0.$$

Thus,

$$y'' = v_1y_1'' + v_1'y_1' + v_2y_2'' + v_2'y_2'.$$

Method of Variation of Parameters

Substituting y, y', y'' in the given non-homogeneous ODE, and rearranging terms, we get:

$$v_1(y_1'' + py_1' + qy_1) + v_2(y_2'' + py_2' + qy_2) + v_1'y_1' + v_2'y_2' = r(x).$$

Thus,

$$v_1'y_1' + v_2'y_2' = r(x).$$

Recall that we also have

$$v_1'y_1 + v_2'y_2 = 0.$$

Thus, we have:

$$\begin{bmatrix} y_1 & y_2 \\ y_1' & y_2' \end{bmatrix} \begin{bmatrix} v_1' \\ v_2' \end{bmatrix} = \begin{bmatrix} 0 \\ r(x) \end{bmatrix}.$$

Method of Variation of Parameters

Therefore,

$$v_1' = \frac{\begin{vmatrix} 0 & y_2 \\ r(x) & y_2' \end{vmatrix}}{W(y_1, y_2)}, \quad v_2' = \frac{\begin{vmatrix} y_1 & 0 \\ y_1' & r(x) \end{vmatrix}}{W(y_1, y_2)}.$$

Thus,

$$v_1 = -\int \frac{y_2 r(x)}{W(y_1, y_2)} dx, \quad v_2 = \int \frac{y_1 r(x)}{W(y_1, y_2)} dx.$$

Hence,

$$\begin{aligned} y &= v_1 y_1 + v_2 y_2 \\ &= y_2 \int \frac{y_1 r(x)}{W(y_1, y_2)} dx - y_1 \int \frac{y_2 r(x)}{W(y_1, y_2)} dx. \end{aligned}$$

Method of Variation of Parameters

Example: Find a particular solution of

$$y'' + y = \csc x.$$

Step I: Find a basis of solutions for the associated homogeneous equation

$$y'' + y = 0.$$

The general solution of this is

$$y(x) = c_1 \sin x + c_2 \cos x.$$

Step II: Calculate the Wronskian $W(y_1, y_2)$:

$$W(y_1, y_2) = \begin{vmatrix} \sin x & \cos x \\ \cos x & -\sin x \end{vmatrix} = -1.$$

Now,

$$v_1 = -\int \frac{y_2 r(x)}{W(y_1, y_2)} dx = -\int \frac{\cos x \csc x}{-1} dx = \ln|\sin x|,$$

and

$$v_2 = \int \frac{y_1 r(x)}{W(y_1, y_2)} dx = \int \frac{\sin x \csc x}{-1} dx = -x.$$

Hence, a particular solution is given by

$$y(x) = \sin x \ln|\sin x| - x \cos x.$$

Method of Variation of Parameters

Example: Find a particular solution of

$$y'' + 4y = 3\cos 2t.$$

Recall that via the method of undetermined coefficients, you had to modify the proposed initial solution by multiplying it by t, and you got the answer as $\frac{3}{4}t\sin 2t$.
Now in variation of parameters,

$$y_1 = \cos 2t,\ \ y_2 = \sin 2t,$$

and

$$v_1 = -\int \frac{\sin 2t \cdot 3\cos 2t}{2}dt = -\frac{3}{16}\cos 4t,$$

$$v_2 = \int \frac{\cos 2t \cdot 3\cos 2t}{2}dt = \frac{3}{16}\sin 4t + \frac{3}{4}t.$$

Thus, a particular solution is

$$v_1y_1 + v_2y_2 = -\frac{3}{16}\cos 6t + \frac{3}{4}t\sin 2t.$$

Method of Variation of Parameters

Example: Find a particular solution of

$$y'' + 3y' + 2y = \frac{1}{1+e^x}.$$

Step I: Find a basis of solutions for the associated homogeneous equation

$$y'' + 3y' + 2y = 0.$$

The general solution of the homogeneous DE is

$$y(x) = c_1 e^{-x} + c_2 e^{-2x}.$$

Step II: We look for a particular solution of the non-homogeneous DE

$$y(x) = v_1 e^{-x} + v_2 e^{-2x}.$$

where

$$v_1' e^{-x} + v_2' e^{-2x} = 0$$

$$-v_1' e^{-x} - 2v_2' e^{-x} = \frac{1}{1+e^x}.$$

Solve for v_1 and v_2 ....